arXiv:1401.0391v3 [hep-ph] 15 Nov 2014

Semileptonic decays $B_c^+ \to D_{(s)}^{(*)}(l^+\nu_l, l^+l^-, \nu\bar{\nu})$ in the perturbative QCD approach

Wen-Fei Wang[1*], Xin Yu[1], Cai-Dian Lü[1], and Zhen-Jun Xiao[2]

[1] *Center for Future High Energy Physics, Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, People's Republic of China,*

[2] *Department of Physics and Institute of Theoretical Physics, Nanjing Normal University, Nanjing, Jiangsu 210023, People's Republic of China*

(Dated: November 18, 2014)

Abstract

In this paper we study the semileptonic decays of $B_c^+ \to D_{(s)}^{(*)}(l^+\nu_l, l^+l^-, \nu\bar{\nu})$ (here l stands for e, μ, or τ). After evaluating the $B_c^+ \to (D_{(s)}, D_{(s)}^*)$ transition form factors $F_{0,+,T}(q^2)$ and $V(q^2), A_{0,1,2}(q^2), T_{1,2,3}(q^2)$ by employing the perturbative QCD factorization approach, we calculate the branching ratios for all these semileptonic decays. Our predictions for the values of the $B_c^+ \to D_{(s)}$ and $B_c^+ \to D_{(s)}^*$ transition form factors are consistent with those obtained by using other methods. The branching ratios of the decay modes with $\bar{\nu}\nu$ are almost an order of magnitude larger than the corresponding decays with l^+l^- after the summation over the three neutrino generations. The branching ratios for the decays with $b \to d$ transitions are much smaller than those decays with the $b \to s$ transitions, due to the Cabibbo-Kobayashi-Maskawa suppression. We define ratios R_D and R_{D^*} for the branching ratios with the τ lepton versus μ, e lepton final states to cancel the uncertainties of the form factors, which could possibly be tested in the near future.

PACS numbers: 13.20.He, 12.38.Bx, 14.40.Nd

* wangwf@ihep.ac.cn

I. INTRODUCTION

The B_c meson is a pseudoscalar ground state of b and c quarks, and thus the electromagnetic interaction cannot transform the B_c meson into other hadrons containing b and c quarks. The two difference of quark flavors forbid its annihilation into gluons and being below the $B-D$ threshold makes the B_c meson stable for strong interaction. The B_c meson can only decay through weak interactions, so it is an ideal system to study weak decays of heavy quarks. Either the heavy quark (b or c) can decay individually, which makes it different from the $B_{u,d}$ or B_s meson. The phase space in the $c \to s$ transition is smaller than that in the $b \to c$ transition, but the Cabibbo-Kobayashi-Maskawa (CKM) matrix element $|V_{cs}| \sim 1$ is much larger than the CKM matrix element $|V_{cb}| \sim 0.04$. Thus the c-quark decays provide the dominant contribution (about 70%) to the decay width of the B_c meson [1]. Because the mass of a $B_c\bar{B}_c$ pair exceeds the threshold of $\Upsilon(4S)$, the B_c meson cannot be produced at the B factories. So comparing with $B_{u,d}$ or B_s meson, the B_c meson decays received much less experimental attention in the past decades. However, at LHC experiments, around 5×10^{10} B_c events per year are expected [1, 2] due to the relatively large production cross section, which provides a very good platform to study various B_c meson decay modes.

Because there is only one hadronic final product, the B_c meson semileptonic decays among the abundant decay modes are relatively clean in the theoretical treatment. These semileptonic decays provide good opportunities to measure not only the CKM matrix elements, such as $|V_{cb}|, |V_{ub}|$, and $|V_{cd}|$, but also the form factors of the B_c to bottom and charmed mesons transitions. The rare semileptonic decays governed by the flavor-changing neutral currents are forbidden at tree level in the standard model (SM). Those decays, which are very sensitive to the contributions of new intermediate particles or interactions are especially interesting. There are various approaches working on the semileptonic B_c decays. In Ref. [3], for example, Dhir and Verma presented a detailed analysis of the exclusive semileptonic B_c decays in the Bauer-Stech-Wirbel framework. The authors of the Refs. [4–6] studied the semileptonic B_c decays in the relativistic and/or constituent quark model. In Refs. [7, 8], $B_c \to D_s^* l^+ l^-$ decays were studied in the SM with the fourth-generation and supersymmetric models. The three point QCD sum rules approach was adopted to investigate the $B_c^+ \to D_{(s)}^{*+} l^+ l^-$ in [9] and $B_c^+ \to D_{(s)}^+ (l^+ l^-, \bar{\nu}\nu)$ in [10].

In this paper, we will study the semileptonic decays of $B_c^+ \to D_{(s)}^{(*)}(l^+\nu_l, l^+l^-, \nu\bar{\nu})$ (here l stands for leptons e, μ, or τ) the perturbative QCD (pQCD) approach [11]. These semileptonic decays are governed by the form factors. At the maximum recoil region, the final state meson is collinear with a large momentum. he spectator c quark in B_c meson thus needs a hard gluon to kick it off from almost zero momentum to a collinear state. However, when doing integrations of momentum fractions of valence quarks, endpoint singularity occurs. A natural way to kill this singularity is to pick up the neglected transverse momentum in the collinear factorization. With the additional transverse momentum cale k_T [12], double logarithms appear in the calculation. We have to use the renormalization group equation to perform the resummation resulting in the so-called Sudakov form factors [13] and make the perturbative calculation of the hard amplitudes (form factors) infrared safe. The pQCD approach is widely adopted to calculate the transition form factors of $B_{u,d}$ and B_s meson[14–16]. Furthermore, various B_c decay modes have also been studied in Refs. [17, 18] in the pQCD approach.

The structure of this paper is as follows. After this Introduction, we collect the distribu-



FIG. 1: The leading-order Feynman diagrams for the transition of $B_c^+ \to (D^{(*)}, D_s^{(*)})$, where M stands for a $D^{(*)}$ or $D_s^{(*)}$ meson, and $\otimes$ is the weak vertex.

tion amplitudes of the B_c, $D^{(*)}$ and $D_s^{(*)}$ mesons in Sec. II. Based on the $k_{\rm T}$ factorization formalism, we calculate and present the expressions for the $B_c \to (D^{(*)}, D_s^{(*)})$ transition form factors in the large recoil regions in Section III. The numerical results and relevant discussions are given in Sec. IV. And Sec. V contains a short summary.

II. KINEMATICS AND THE WAVE FUNCTIONS

The lowest-order diagrams for $B_c^+ \to (D^{(*)}, D_s^{(*)})$ transitions are displayed in Fig. 1, where M stands for $D^{(*)}$ or $D_s^{(*)}$ meson, the $\otimes$ is the weak vertex for the leptonic pairs to come out. In the rest frame of B_c meson, with the m_{B_c} standing for the mass of the B_c meson, and m for the $D_{(s)}$ or $D^*_{(s)}$ mesons, the momenta of B_c and $D^{(*)}_{(s)}$ mesons are defined in the light-cone coordinates as [18, 19]

$$p_1 = \frac{m_{B_c}}{\sqrt{2}}(1, 1, 0_\perp), \quad p_2 = \frac{m_{B_c}}{\sqrt{2}}(r\eta^+, r\eta^-, 0_\perp), \tag{1}$$

with $r = m/m_{B_c}$ and $\eta^\pm = \eta \pm \sqrt{\eta^2 - 1}$. As for the η in $\eta^\pm$, the expression

$$\eta = \frac{1}{2r}\left[1 + r^2 - \frac{q^2}{m_{B_c}^2}\right] \tag{2}$$

can be evaluated from $q^2 = (p_1 - p_2)^2$ which is the invariant mass of the lepton pairs. The momenta of the spectator quarks in the B_c and $D^{(*)}_{(s)}$ mesons are parameterized as

$$k_1 = (x_1\frac{m_{B_c}}{\sqrt{2}}, x_1\frac{m_{B_c}}{\sqrt{2}}, k_{1\perp}), \quad k_2 = (x_2\frac{m_{B_c}}{\sqrt{2}}r\eta^+, x_2\frac{m_{B_c}}{\sqrt{2}}r\eta^-, k_{2\perp}). \tag{3}$$

For the $D^*_{(s)}$ mesons, we define their polarization vector ϵ as

$$\epsilon_L = \frac{1}{\sqrt{2}}(\eta^+, -\eta^-, 0_\perp), \quad \epsilon_T = (0, 0, 1), \tag{4}$$

where ϵ_L and ϵ_T denote the longitudinal and transverse polarization of the $D^*_{(s)}$ mesons, respectively.

In this work, we use the same distribution amplitude for the B_c meson as that used in Refs. [18, 20–22],

$$\Phi_{B_c}(x) = \frac{i}{\sqrt{2N_c}}[(\not{p} + m_{B_c})\gamma_5\phi_{B_c}(x)]_{\alpha\beta}\ , \tag{5}$$

with

$$\phi_{B_c}(x) = \frac{f_{B_c}}{2\sqrt{2N_c}}\delta(x - m_c/m_{B_c})\exp[-\omega_{B_c}^2 b^2/2]\ , \tag{6}$$

where m_c is the mass of c-quark. Because the B_c meson consists of two heavy quarks b and c, just like a heavy quarkonium, the non-relativistic QCD framework can be applied, which means the leading-order wave function should be just the zero-point wave function shown in Eq. (6).

For the $D_{(s)}^{(*)}$ mesons, up to twist-3 accuracy, the two-parton light-cone distribution amplitudes are defined as [19, 23]

$$\begin{aligned}
\langle D_{(s)}(p)|q_\alpha(z)\bar{c}_\beta(0)|0\rangle &= \frac{i}{\sqrt{2N_C}}\int_0^1 dx e^{ixp\cdot z}\left[\gamma_5\left(\not{p}+m\right)\phi_{D_{(s)}}(x,b)\right]_{\alpha\beta}\,, \\
\langle D_{(s)}^*(p)|q_\alpha(z)\bar{c}_\beta(0)|0\rangle &= -\frac{1}{\sqrt{2N_C}}\int_0^1 dx e^{ixp\cdot z}\left[\not{\epsilon}_L(\not{p}+m)\phi_{D_{(s)}^*}(x,b)\right. \\
&\quad \left.+\not{\epsilon}_T(\not{p}+m)\phi_{D_{(s)}^*}(x,b)\right]_{\alpha\beta}\,,
\end{aligned} \tag{7}$$

with

$$\begin{aligned}
\int_0^1 dx\phi_{D_{(s)}}(x,0) &= \frac{f_{D_{(s)}}}{2\sqrt{2N_c}}\,, \\
\int_0^1 dx\phi_{D_{(s)}^*}(x,0) &= \frac{f_{D_{(s)}^*}}{2\sqrt{2N_c}}\,,
\end{aligned} \tag{8}$$

as the normalization conditions. We adopt $f_D = 206.7\pm8.9$ MeV and $f_{D_s} = 260.0\pm5.6$ MeV in PDG [24] by experimental average for D and D_s mesons, respectively. For the D^* or D_s^* meson, we adopt the same decay constant and distribution amplitude for the longitudinal and transverse components. Since there is no experimental data, we use $f_{D^*} = 270$ MeV and $f_{D_s^*} = 310$ MeV for D^* and D_s^* meson considering of the results in Refs. [25] and assume a 10% uncertainty. The distribution amplitude for the $D_{(s)}$ meson is

$$\phi_{D_{(s)}} = \frac{1}{2\sqrt{2N_C}}f_{D_{(s)}}6x(1-x)\left[1+C_{D_{(s)}}(1-2x)\right]\exp\left[-\frac{\omega_{D_{(s)}}^2 b^2}{2}\right], \tag{9}$$

which is a k_T-dependent form with $C_D = 0.5, \omega_D = 0.1$ and $C_{D_s} = 0.4, \omega_{D_s} = 0.2$ for D and D_s mesons, respectively [23]. In this work, we also adopt the same distribution amplitude for both the vector meson $D_{(s)}^*$ and pseudoscalar meson $D_{(s)}$ because of their small mass difference [23].

III. FORM FACTORS OF SEMILEPTONIC DECAYS

The form factors $F_+(q^2), F_0(q^2)$ for the B_c to pseudoscalar meson $D_{(s)}$ transition induced by the vector current can be defined as [26, 27]

$$\begin{aligned}
\langle D_{(s)}(p_2)|\bar{q}(0)\gamma_\mu b(0)|B_c(p_1)\rangle &= \left[(p_1+p_2)_\mu - \frac{m_{B_c}^2-m^2}{q^2}q_\mu\right]F_+(q^2) \\
&+ \frac{m_{B_c}^2-m^2}{q^2}q_\mu F_0(q^2),
\end{aligned} \tag{10}$$

where $q = p_1 - p_2$ is the momentum of the lepton pairs. In order to cancel the poles at $q^2 = 0$, $F_+(0)$ should be equal to $F_0(0)$. For the sake of convenience, we define the auxiliary form factors $f_1(q^2)$ and $f_2(q^2)$,

$$\langle D_{(s)}(p_2)|\bar{q}(0)\gamma_\mu b(0)|B_c(p_1)\rangle = f_1(q^2)p_{1\mu} + f_2(q^2)p_{2\mu}. \tag{11}$$

In terms of $f_1(q^2)$ and $f_2(q^2)$ the form factors $F_+(q^2)$ and $F_0(q^2)$ are

$$\begin{aligned} F_+(q^2) &= \frac{1}{2}\left[f_1(q^2) + f_2(q^2)\right], \\ F_0(q^2) &= \frac{1}{2}f_1(q^2)\left[1 + \frac{q^2}{m_{B_c}^2 - m^2}\right] + \frac{1}{2}f_2(q^2)\left[1 - \frac{q^2}{m_{B_c}^2 - m^2}\right]. \end{aligned} \tag{12}$$

The form factor $F_T(q^2)$ for the $B_c \to D_{(s)}$ transition induced by the tensor current can be defined as [27]

$$\langle D_{(s)}(p_2)|\bar{q}(0)\sigma_{\mu\nu}b(0)|B_c(p_1)\rangle = i\left[p_{2\mu}q_\nu - q_\mu p_{2\nu}\right]\frac{2F_T(q^2)}{m_{B_c} + m}\,. \tag{13}$$

There are seven form factors $V(q^2), A_{0,1,2}(q^2)$ and $T_{1,2,3}(q^2)$ needed for the transition of $B_c \to D^*_{(s)}$ in this work. The form factors $V(q^2)$ and $A_{0,1,2}(q^2)$ are defined by [27–29]

$$\langle D^*_{(s)}(p_2)|\bar{q}(0)\gamma_\mu b(0)|B_c(p_1)\rangle = \epsilon_{\mu\nu\alpha\beta}\epsilon^{\nu *}p_1^\alpha p_2^\beta \frac{2V(q^2)}{m_{B_c} + m}\,, \tag{14}$$

$$\begin{aligned} \langle D^*_{(s)}(p_2)|\bar{q}(0)\gamma_\mu\gamma_5 b(0)|B_c(p_1)\rangle &= i\left[\epsilon^*_\mu - \frac{\epsilon^*\cdot q}{q^2}q_\mu\right](m_{B_c} + m)A_1(q^2) \\ &- i\left[(p_1 + p_2)_\mu - \frac{m_{B_c}^2 - m^2}{q^2}q_\mu\right](\epsilon^*\cdot q)\frac{A_2(q^2)}{m_{B_c} + m} \\ &+ i\frac{2m(\epsilon^*\cdot q)}{q^2}q_\mu A_0(q^2), \end{aligned} \tag{15}$$

where ϵ^* is the polarization vector of the $D^*_{(s)}$ meson. The form factors $T_{1,2,3}$ are defined by [27, 30]

$$\begin{aligned} \langle D^*_{(s)}(p_2)|\bar{q}(0)\sigma_{\mu\nu}q^\nu(1 + \gamma_5)b(0)|B_c(p_1)\rangle &= i\epsilon_{\mu\nu\alpha\beta}\epsilon^{*\nu}p_1^\alpha p_2^\beta 2T_1(q^2) \\ &+ \left[\epsilon^*_\mu(m_{B_c}^2 - m^2) - (\epsilon^*\cdot q)(p_1 + p_2)_\mu\right]T_2(q^2) \\ &+ (\epsilon^*\cdot q)\left[q_\mu - \frac{q^2}{m_{B_c}^2 - m^2}(p_1 + p_2)_\mu\right]T_3(q^2)\,, \end{aligned} \tag{16}$$

with $T_1(0) = T_2(0)$ implied by the identity

$$\sigma_{\mu\nu}\gamma_5 = -\frac{i}{2}\epsilon_{\mu\nu\alpha\beta}\sigma^{\alpha\beta}\,. \tag{17}$$

In the transverse configuration b-space and by including the Sudakov form factors and the threshold resummation effects, we obtain the $B_c \to D_{(s)}$ form factors $f_1(q^2), f_2(q^2)$ and

$F_T(q^2)$ as follows

$$
\begin{aligned}
f_1(q^2) &= 16\pi m_{B_c}^2 r C_F \int dx_1 dx_2 \int b_1 db_1 b_2 db_2 \; \phi_{B_c}(x_1)\phi_{D_{(s)}}(x_2,b_2) \\
&\times \Big\{ [1-rx_2]\cdot h_1(x_1,x_2,b_1,b_2)\cdot \alpha_s(t_1)\exp\left[-S_{ab}(t_1)\right] \\
&- \; [r+2x_1(1-\eta)]\cdot h_2(x_1,x_2,b_1,b_2)\cdot \alpha_s(t_2)\exp\left[-S_{ab}(t_2)\right] \Big\} \,, \qquad (18)
\end{aligned}
$$

$$
\begin{aligned}
f_2(q^2) &= 16\pi m_{B_c}^2 C_F \int dx_1 dx_2 \int b_1 db_1 b_2 db_2 \; \phi_{B_c}(x_1)\phi_{D_{(s)}}(x_2,b_2) \\
&\times \Big\{ [1-2rx_2(1-\eta)]\cdot h_1(x_1,x_2,b_1,b_2)\cdot \alpha_s(t_1)\exp\left[-S_{ab}(t_1)\right] \\
&+ \; [2r-x_1]\cdot h_2(x_1,x_2,b_1,b_2)\cdot \alpha_s(t_2)\exp\left[-S_{ab}(t_2)\right] \Big\} \,, \qquad (19)
\end{aligned}
$$

$$
\begin{aligned}
F_T(q^2) &= 8\pi m_{B_c}^2 C_F (1+r) \int dx_1 dx_2 \int b_1 db_1 b_2 db_2 \; \phi_{B_c}(x_1)\phi_{D_{(s)}}(x_2,b_2) \\
&\times \Big\{ [1-rx_2]\cdot h_1(x_1,x_2,b_1,b_2)\cdot \alpha_s(t_1)\exp\left[-S_{ab}(t_1)\right] \\
&+ \; [2r-x_1]\cdot h_2(x_1,x_2,b_1,b_2)\cdot \alpha_s(t_2)\exp\left[-S_{ab}(t_2)\right] \Big\} \,, \qquad (20)
\end{aligned}
$$

where $C_F = 4/3$ is the color factor. The functions h_1 and h_2, the scales t_1, t_2 and the Sudakov factors S_{ab} are the same as those given in Refs. [18, 19].

The expressions of form factors $V(q^2)$, $A_{0,1,2}(q^2)$ and $T_{1,2,3}(q^2)$ for the $B_c \to D_{(s)}^*$ transition in the pQCD approach are:

$$
\begin{aligned}
V(q^2) &= 8\pi m_{B_c}^2 C_F (1+r) \int dx_1 dx_2 \int b_1 db_1 b_2 db_2 \; \phi_{B_c}(x_1)\phi^T_{D^*_{(s)}}(x_2,b_2) \\
&\times \Big\{ [1-rx_2]\cdot h_1(x_1,x_2,b_1,b_2)\cdot \alpha_s(t_1)\exp\left[-S_{ab}(t_1)\right] \\
&+ \; r\cdot h_2(x_1,x_2,b_1,b_2)\cdot \alpha_s(t_2)\exp\left[-S_{ab}(t_2)\right] \Big\} \,, \qquad (21)
\end{aligned}
$$

$$
\begin{aligned}
A_0(q^2) &= 8\pi m_{B_c}^2 C_F \int dx_1 dx_2 \int b_1 db_1 b_2 db_2 \; \phi_{B_c}(x_1)\phi^L_{D^*_{(s)}}(x_2,b_2) \\
&\times \Big\{ [1-rx_2(r-2\eta)+r\,(1-2x_2)] \\
&\times \; h_1(x_1,x_2,b_1,b_2)\cdot \alpha_s(t_1)\exp\left[-S_{ab}(t_1)\right] \\
&+ \; \left[r^2+x_1(1-2r\eta)\right]\cdot h_2(x_1,x_2,b_1,b_2)\cdot \alpha_s(t_2)\exp\left[-S_{ab}(t_2)\right] \Big\} \,, \qquad (22)
\end{aligned}
$$

$$
\begin{aligned}
A_1(q^2) &= 16\pi m_{B_c}^2 C_F \frac{r}{1+r} \int dx_1 dx_2 \int b_1 db_1 b_2 db_2 \; \phi_{B_c}(x_1)\phi^T_{D^*_{(s)}}(x_2,b_2) \\
&\times \Big\{ [1+rx_2\eta-2rx_2+\eta]\cdot h_1(x_1,x_2,b_1,b_2)\cdot \alpha_s(t_1)\exp[-S_{ab}(t_1)] \\
&+ \; [r\eta-x_1]\cdot h_2(x_1,x_2,b_1,b_2)\cdot \alpha_s(t_2)\exp[-S_{ab}(t_2)] \Big\} \,, \qquad (23)
\end{aligned}
$$

$$
\begin{aligned}
A_2(q^2) &= \frac{(1+r)^2(\eta-r)}{2r(\eta^2-1)}\cdot A_1(q^2) - 8\pi m_{B_c}^2 C_F \frac{1+r}{\eta^2-1}\int dx_1 dx_2 \int b_1 db_1 b_2 db_2\ \phi_{B_c}(x_1) \\
&\times\ \phi^L_{D^*_{(s)}}(x_2,b_2)\cdot\Big\{\big[\eta(1-r^2x_2) - rx_2(1-2\eta^2-2r) + (1-r) - r\eta(1+2x_2)\big] \\
&\times\ h_1(x_1,x_2,b_1,b_2)\cdot\alpha_s(t_1)\exp\left[-S_{ab}(t_1)\right] \\
&+\ \big[r(1-x_1+2x_1\eta^2) - \eta(r^2+x_1)\big] \\
&\times\ h_2(x_1,x_2,b_1,b_2)\cdot\alpha_s(t_2)\exp\left[-S_{ab}(t_2)\right]\Big\}\,, \qquad (24)
\end{aligned}
$$

$$
\begin{aligned}
T_1(q^2) &= 8\pi m_{B_c}^2 C_F \int dx_1 dx_2 \int b_1 db_1 b_2 db_2\ \phi_{B_c}(x_1)\phi^T_{D^*_{(s)}}(x_2,b_2) \\
&\times\ \Big\{[1+r(1-x_2(2+r-2\eta))]\cdot h_1(x_1,x_2,b_1,b_2)\cdot\alpha_s(t_1)\exp\left[-S_{ab}(t_1)\right] \\
&+\ r\left[1-x_1\right]\cdot h_2(x_1,x_2,b_1,b_2)\cdot\alpha_s(t_2)\exp\left[-S_{ab}(t_2)\right]\Big\}\,, \qquad (25)
\end{aligned}
$$

$$
\begin{aligned}
T_2(q^2) &= 16\pi m_{B_c}^2 C_F \frac{r}{1-r^2}\int dx_1 dx_2 \int b_1 db_1 b_2 db_2\ \phi_{B_c}(x_1)\phi^T_{D^*_{(s)}}(x_2,b_2) \\
&\times\ \Big\{\big[(1-r)(1+\eta) + 2rx_2(r-\eta) + rx_2(2\eta^2 - r\eta - 1)\big] \\
&\times\ h_1(x_1,x_2,b_1,b_2)\cdot\alpha_s(t_1)\exp\left[-S_{ab}(t_1)\right] \\
&+\ \big[r(1+x_1)\eta - r^2 - x_1\big]\cdot h_2(x_1,x_2,b_1,b_2)\cdot\alpha_s(t_2)\exp\left[-S_{ab}(t_2)\right]\Big\}\,, \qquad (26)
\end{aligned}
$$

$$
\begin{aligned}
T_3(q^2) &= \frac{r+\eta}{r}\cdot\frac{1-r^2}{2(\eta^2-1)}\cdot T_2(q^2) - \frac{1-r^2}{(\eta^2-1)} \\
&\times\ 8\pi m_{B_c}^2 C_F \int dx_1 dx_2 \int b_1 db_1 b_2 db_2\ \phi_{B_c}(x_1)\phi^L_{D^*_{(s)}}(x_2,b_2) \\
&\times\ \Big\{[1+rx_2(\eta-2)+\eta]\cdot h_1(x_1,x_2,b_1,b_2)\cdot\alpha_s(t_1)\exp\left[-S_{ab}(t_1)\right] \\
&+\ [x_1\eta - r]\cdot h_2(x_1,x_2,b_1,b_2)\cdot\alpha_s(t_2)\exp\left[-S_{ab}(t_2)\right]\Big\}\,. \qquad (27)
\end{aligned}
$$

One should note that the expressions for the form factors $f_{1,2}(q^2)$, $F_T(q^2)$, $V(q^2)$, $A_{0,1,2}(q^2)$ and $T_{1,2,3}(q^2)$ given in Eqs. (18)-(27) are the results at leading order of the pQCD approach. The next-to-leading-order contributions to the form factors of $B\to(\pi,K,\eta^{(\prime)})$ transitions given in Refs. [14, 15, 31] are not available here because of the large mass of c-quark and $(D_{(s)}, D^*_{(s)})$ mesons.

One should note that the pQCD predictions for the considered form factors are reliable only for the small values of q^2. For the form factors in the large-q^2 region, one has to make an extrapolation for them from the low-q^2 region to large-q^2 region. In this work we make the extrapolation by using the formula in Refs. [18, 32]

$$
F(q^2) = F(0)\cdot\exp\left[a\cdot q^2 + b\cdot(q^2)^2\right], \qquad (28)
$$

where F stands for the form factors $F_{0,+,T}, V, A_{0,1,2}$ and $T_{1,2,3}$, and a, b are the constants to be determined by the fitting procedure.

The $B_c^- \to \bar{D}^0 l^- \bar{\nu}_l$ and $B_c^- \to \bar{D}^{*0} l^- \bar{\nu}_l$ decays are from the quark level $b \to u l^- \bar{\nu}$ charged current transition. The effective Hamiltonian for such transition is [33]

$$\mathcal{H}_{eff}(b \to u l^- \bar{\nu}_l) = \frac{G_F}{\sqrt{2}} V_{ub}\, \bar{u}\gamma_\mu(1-\gamma_5) b \cdot \bar{l}\gamma^\mu(1-\gamma_5)\nu_l, \tag{29}$$

where $G_F = 1.16637 \times 10^{-5} GeV^{-2}$ is the Fermi-coupling constant and V_{ub} is one of the CKM matrix elements. With the form factors calculated in Eqs. (18,19,21-24), one can easily get the differential decay width expression for $B_c^- \to \bar{D}^0 l^- \bar{\nu}_l$ and $B_c^- \to \bar{D}^{*0} l^- \bar{\nu}_l$.

For those flavor-changing neutral-current one-loop decay modes, such as $B_c \to D^{(*)} l^+ l^-$ and $B_c \to D_s^{(*)} l^+ l^-$, are transitions of $b \to d l^+ l^-$ and $b \to s l^+ l^-$ at quark level, respectively. The effective Hamiltonians and the corresponding differential decay widths are more complicated, we refer the readers to Refs. [14, 30, 34–36].

For the decay modes of $B_c \to D_s^{(*)} \nu\bar{\nu}$, the effective Hamiltonian is [33]

$$\mathcal{H}_{eff}(b \to s\nu\bar{\nu}) \;=\; \frac{G_F}{\sqrt{2}} \frac{\alpha_{em}}{2\pi \sin^2(\theta_W)} V_{tb} V_{ts}^* \eta_X X(x_t)\; [\bar{s}\gamma^\mu(1-\gamma_5) b]\, [\bar{\nu}\gamma_\mu(1-\gamma_5)\nu] \tag{30}$$

where θ_W is the Weinberg angle with $\sin^2(\theta_W) = 0.231$ [24], V_{tb} and V_{ts} are CKM matrix elements and $\alpha_{em} \approx 1/137$ is the fine structure constant. The function $X(x_t)$ can be found in Ref. [33], while $\eta_X \approx 1$ is the QCD radiative correction factor [33]. As for the decay modes of $B_c \to D^{(*)} \nu\bar{\nu}$, their effective Hamiltonian can be obtained by a simple replacement of $s \to d$ in Eq. (30). The corresponding differential decay widths for $B_c \to D_{(s)} \nu\bar{\nu}$ is the same as $B \to \pi(K) \nu\bar{\nu}$ in Ref. [14] except the replacements $m_B \to m_{B_c}$ and $m_P \to m$. While for the decay modes of $B_c \to D_s^* \nu\bar{\nu}$, the differential decay width is [37]

$$\begin{aligned}
\frac{d\Gamma(B_c \to D_s^* \nu\bar{\nu})}{dq^2} \;=\; & \frac{G_F^2 \alpha_{em}^2}{2^{10}\pi^5 m_{B_c}^3} \cdot \left| \frac{X(x_t)}{sin^2(\theta_w)} \right|^2 \cdot \eta_X^2 \cdot |V_{tb} V_{ts}^*|^2\, \lambda^{\frac{1}{2}} \Bigg\{ 8\lambda q^2 \frac{V^2}{(m_{B_c}+m)^2} \\
& + \frac{1}{m^2} \Bigg[\lambda^2 \frac{A_2^2}{(m_{B_c}+m)^2} + (m_{B_c}+m)^2 (\lambda + 12 m^2 q^2) \cdot A_1^2 \\
& - 2\lambda (m_{B_c}^2 - m^2 - q^2) \cdot Re[A_1^* A_2] \Bigg] \Bigg\}\,,
\end{aligned} \tag{31}$$

where V, A_1 and A_2 are the form factors of $B_c \to D_s^*$ transition, and the phase-space factor

$$\lambda = (m_{B_c}^2 + m^2 - q^2)^2 - 4 m_{B_c}^2 m^2\,. \tag{32}$$

IV. NUMERICAL RESULTS AND DISCUSSIONS

In the numerical calculations we adopt the following input parameters [24]

$$\begin{aligned}
m_{B_c^-} \;&=\; 6.277 \text{ GeV}, \quad m_{\bar{D}^0} = 1.865 \text{ GeV}, \quad m_{D^-} = 1.870 \text{ GeV}, \\
m_{\bar{D}^{*0}} \;&=\; 2.007 \text{ GeV}, \quad m_{D^{*-}} = 2.010 \text{ GeV}, \quad m_{D_s^-} = 1.969 \text{ GeV}, \\
m_{\bar{D}_s^{*-}} \;&=\; 2.112 \text{ GeV}, \quad m_\tau = 1.777 \text{ GeV}, \quad m_c = 1.275 \pm 0.025 \text{ GeV}, \\
\tau_{B_c} \;&=\; (0.45 \pm 0.04) \text{ ps},
\end{aligned} \tag{33}$$

For the CKM matrix element V_{ub}, we adopt the value in Refs. [14, 38]. And we use $|V_{tb}| = 0.999, |V_{ts}| = 0.0404$ and $|V_{td}/V_{ts}| = 0.211$ [24] in this work. As for the decay constant of

the B_c meson, we adopt 0.489 GeV [39] as its central value, and give it an uncertainty of 0.050 GeV.

The numerical values of the $B_c \to D$ and $B_c \to D_s$ transition form factors $F_{0,+,T}$ at $q^2 = 0$ and their fitted parameters a, b are listed in Table I. The numerical values of the form factors $V, A_{0,1,2}$ and $T_{1,2,3}$ at $q^2 = 0$ for the $B_c \to D^*$ and $B_c \to D_s^*$ transitions are listed in Table II. The first error of the pQCD predictions for the form factors in Table I and Table II is induced by the B_c meson wave function parameter $\omega_{B_c} = 1.0 \pm 0.1$; the second error comes from the uncertainty of decay constant f_{B_c}; the third error comes from the uncertainty of decay constants of the $D_{(s)}^{(*)}$ mesons; the fourth error in Tables I and II comes from the uncertainty of $D_{(s)}^{(*)}$ wave function $C_{D^{(*)}} = 0.5 \pm 0.1$ or $C_{D_s^{(*)}} = 0.4 \pm 0.1$; the fifth error comes from $m_c = 1.275 \pm 0.025$ GeV. The errors from the uncertainty of $\omega_{D^{(*)}} = 0.10 \pm 0.02$ or $\omega_{D_s^{(*)}} = 0.20 \pm 0.04$ are very small that have been neglected.

Unlike the form factors at maximum recoil, the extrapolation parameters a, b of the form factors are less sensitive to the decay constant and wave function of $D_{(s)}^{(*)}$ meson. In Tables I and II, we only show uncertainties for the parameter a and b from B_c meson wave function parameter ω_{B_c}, and from quark mass uncertainty $m_c = 1.275 \pm 0.025$ GeV. As a comparison, we also present some results obtained by other authors based on different methods in Table III. It is easy to see that our results are consistent with the results in literature.

TABLE I: The pQCD predictions for form factors F_0, F_+ and F_T at $q^2 = 0$ and the parametrization constants a and b for $B_c \to D$ and $B_c \to D_s$ transitions.

	$F(0)$	a	b
$F_0^{B_c\to D}$	$0.19 \pm 0.02 \pm 0.02 \pm 0.01 \pm 0.01 \pm 0.01$	$0.038 \pm 0.001 \pm 0.000$	0.0013 ± 0.0001
$F_+^{B_c\to D}$	$0.19 \pm 0.02 \pm 0.02 \pm 0.01 \pm 0.01 \pm 0.01$	$0.059 \pm 0.001 \pm 0.001$	0.0020 ± 0.0001
$F_T^{B_c\to D}$	$0.20 \pm 0.02 \pm 0.02 \pm 0.01 \pm 0.01 \pm 0.01$	$0.070 \pm 0.001 \pm 0.001$	$0.0021^{+0.0000}_{-0.0001}$
$F_0^{B_c\to D_s}$	$0.27 \pm 0.03 \pm 0.03 \pm 0.02 \pm 0.01 \pm 0.01$	$0.039 \pm 0.002 \pm 0.001$	$0.0015^{+0.0001}_{-0.0000}$
$F_+^{B_c\to D_s}$	$0.27 \pm 0.03 \pm 0.03 \pm 0.02 \pm 0.01 \pm 0.01$	$0.061 \pm 0.002 \pm 0.001$	$0.0023^{+0.0001}_{-0.0000}$
$F_T^{B_c\to D_s}$	$0.28 \pm 0.03 \pm 0.03 \pm 0.02 \pm 0.01 \pm 0.01$	$0.073 \pm 0.002 \pm 0.001$	$0.0025^{+0.0000}_{-0.0001}$

With the form factors given, it is straightforward to calculate the branching ratios for all the considered semileptonic decays by performing the numerical integration over the whole range of q^2. For the $b \to u$ charged current process, with $l = (e, \mu)$, the decay rates are the following:

$$\begin{aligned}
Br(B_c^- \to \bar{D}^0 l^- \bar{\nu}_l) &= (3.15^{+0.97}_{-0.72}(\omega_{B_c})^{+0.68}_{-0.61}(f_{B_c})^{+0.29}_{-0.27}(m_c)^{+0.31}_{-0.29}(C_D)^{+0.28}_{-0.27}(f_D) \pm 0.28(\tau_{B_c})) \cdot 10^{-5}, \\
Br(B_c^- \to \bar{D}^0 \tau^- \bar{\nu}_\tau) &= (2.16^{+0.72}_{-0.52}(\omega_{B_c})^{+0.46}_{-0.42}(f_{B_c})^{+0.22}_{-0.19}(m_c)^{+0.20}_{-0.19}(C_D)^{+0.19}_{-0.18}(f_D) \pm 0.19(\tau_{B_c})) \cdot 10^{-5}, \\
Br(B_c^- \to \bar{D}^{*0} l^- \bar{\nu}_l) &= (1.09^{+0.34}_{-0.26}(\omega_{B_c})^{+0.23}_{-0.21}(f_{B_c})^{+0.13}_{-0.11}(m_c) \pm 0.10(C_{D^*})^{+0.23}_{-0.21}(f_{D^*}) \pm 0.10(\tau_{B_c})) \cdot 10^{-4}, \\
Br(B_c^- \to \bar{D}^{*0} \tau^- \bar{\nu}_\tau) &= (0.64^{+0.20}_{-0.16}(\omega_{B_c})^{+0.14}_{-0.12}(f_{B_c})^{+0.08}_{-0.07}(m_c)^{+0.06}_{-0.05}(C_{D^*})^{+0.13}_{-0.12}(f_{D^*}) \pm 0.06(\tau_{B_c})) \cdot 10^{-4},
\end{aligned} \tag{34}$$

where the errors come from the uncertainties of $\omega_{B_c} = 1.0 \pm 0.1$, $f_{B_c} = 0.489 \pm 0.050$ GeV, $m_c = (1.275 \pm 0.025)$ GeV, $C_{D^{(*)}} = 0.5 \pm 0.1$, $f_D = (206.7 \pm 8.9)$ MeV or $f_{D^*} = (270 \pm 27)$ MeV and $\tau_{B_c} = (0.45 \pm 0.04)$ ps, respectively.

For the flavor-changing neutral-current processes, after making the numerical integration over the whole range of $4m_l^2 \le q^2 \le (m_{B_c} - m)^2$, we get the pQCD predictions for the

TABLE II: The pQCD predictions for form factors $A_{0,1,2}, V$ and $T_{1,2,3}$ at $q^2 = 0$ and the parametrization constants a and b for $B_c \to D^*$ and $B_c \to D_s^*$ transitions.

	$F(0)$	a	b
$A_0^{B_c\to D^*}$	$0.17 \pm 0.02 \pm 0.02 \pm 0.02 \pm 0.01 \pm 0.00$	$0.063 \pm 0.001 \pm 0.001$	$0.0024 \pm 0.0000 \pm 0.0000$
$A_1^{B_c\to D^*}$	$0.18 \pm 0.02 \pm 0.02 \pm 0.02 \pm 0.01 \pm 0.01$	$0.043 \pm 0.001 \pm 0.001$	$0.0018 \pm 0.0001 \pm 0.0001$
$A_2^{B_c\to D^*}$	$0.20 \pm 0.02 \pm 0.02 \pm 0.02 \pm 0.01 \pm 0.01$	$0.067 \pm 0.001 \pm 0.001$	$0.0026 \pm 0.0001 \pm 0.0001$
$V^{B_c\to D^*}$	$0.25 \pm 0.03 \pm 0.03 \pm 0.03 \pm 0.01 \pm 0.01$	$0.073 \pm 0.002 \pm 0.001$	$0.0029 \pm 0.0001 \pm 0.0001$
$T_1^{B_c\to D^*}$	$0.22 \pm 0.02 \pm 0.02 \pm 0.02 \pm 0.01 \pm 0.01$	$0.063 \pm 0.001 \pm 0.001$	$0.0027 \pm 0.0001 \pm 0.0001$
$T_2^{B_c\to D^*}$	$0.22 \pm 0.02 \pm 0.02 \pm 0.02 \pm 0.01 \pm 0.01$	$0.038 \pm 0.001 \pm 0.001$	$0.0017 \pm 0.0001 \pm 0.0001$
$T_3^{B_c\to D^*}$	$0.20 \pm 0.02 \pm 0.02 \pm 0.02 \pm 0.01 \pm 0.01$	$0.077 \pm 0.002 \pm 0.001$	$0.0049 \pm 0.0001 \pm 0.0001$
$A_0^{B_c\to D_s^*}$	$0.21 \pm 0.02 \pm 0.02 \pm 0.02 \pm 0.01 \pm 0.01$	$0.064 \pm 0.001 \pm 0.001$	$0.0031 \pm 0.0002 \pm 0.0001$
$A_1^{B_c\to D_s^*}$	$0.23 \pm 0.02 \pm 0.02 \pm 0.02 \pm 0.01 \pm 0.01$	$0.044 \pm 0.002 \pm 0.001$	$0.0022 \pm 0.0002 \pm 0.0001$
$A_2^{B_c\to D_s^*}$	$0.25 \pm 0.03 \pm 0.03 \pm 0.03 \pm 0.01 \pm 0.01$	$0.069 \pm 0.002 \pm 0.001$	$0.0035 \pm 0.0002^{+0.0002}_{-0.0001}$
$V^{B_c\to D_s^*}$	$0.33 \pm 0.03 \pm 0.03 \pm 0.03 \pm 0.02 \pm 0.01$	$0.075 \pm 0.002 \pm 0.001$	$0.0039 \pm 0.0002 \pm 0.0001$
$T_1^{B_c\to D_s^*}$	$0.28 \pm 0.03 \pm 0.03 \pm 0.03 \pm 0.02 \pm 0.01$	$0.064 \pm 0.001 \pm 0.001$	$0.0035 \pm 0.0002 \pm 0.0001$
$T_2^{B_c\to D_s^*}$	$0.28 \pm 0.03 \pm 0.03 \pm 0.03 \pm 0.02 \pm 0.01$	$0.039 \pm 0.001 \pm 0.001$	$0.0023 \pm 0.0002 \pm 0.0001$
$T_3^{B_c\to D_s^*}$	$0.27 \pm 0.03 \pm 0.03 \pm 0.03 \pm 0.02 \pm 0.01$	$0.082 \pm 0.002 \pm 0.001$	$0.0068 \pm 0.0002 \pm 0.0002$

TABLE III: Comparison of $B_c \to D_{(s)}^{(*)}$ transition form factors at $q^2 = 0$ evaluated in this paper with other methods.

$B_c \to D^{(*)}$	$F_+(0) = F_0(0)$	$F_T(0)$	$A_0(0)$	$A_1(0)$	$A_2(0)$	$V(0)$	$T_1(0) = T_2(0)$	$T_3(0)$
pQCD	0.19	0.20	0.17	0.18	0.20	0.25	0.22	0.20
Ref.[32]	0.16	–	0.09	0.08	0.07	0.13	–	–
Ref.[40]	0.14	–	0.14	0.17	0.19	0.18	–	–
Ref.[41]	0.189	–	0.284	0.146	0.158	0.296	–	–
Ref.[42]	0.35	–	0.05	0.32	0.57	0.57	–	–
Ref.[43]	0.32	–	0.35	0.43	0.51	1.66	–	–
Ref.[44]	0.075	–	0.081	0.095	0.11	0.16	–	–
$B_c \to D_s^{(*)}$	$F_+(0) = F_0(0)$	$F_T(0)$	$A_0(0)$	$A_1(0)$	$A_2(0)$	$V(0)$	$T_1(0) = T_2(0)$	$T_3(0)$
pQCD	0.27	0.28	0.21	0.23	0.25	0.33	0.28	0.27
Ref.[32]	0.28	–	0.17	0.14	0.12	0.23	–	–
Ref.[43]	0.45	–	0.47	0.56	0.65	2.02	–	–
Ref.[44]	0.15	–	0.16	0.18	0.20	0.29	–	–

branching ratios of considered decay modes which are listed in Table IV. The errors of the pQCD predictions in Table IV come from the uncertainties of ω_{B_c}, m_c, $C_{D^{(*)}}$ or $C_{D_s^{(*)}}$, $f_{D^{(*)}}$ or $f_{D_s^{(*)}}$ and τ_{B_c}, respectively.

From the pQCD predictions for the form factors $F_{0,+,T}$ in Table I, the form factors $V, A_{0,1,2}$ and $T_{1,2,3}$ in Table II and the pQCD predictions for the branching ratios as listed in Eq. (34)

TABLE IV: The pQCD predictions for the branching ratios of the considered decays ($l = e, \mu$).

Decay modes	pQCD predictions
$Br(B_c^- \to D^- l^+ l^-)$	$(3.79^{+1.16}_{-0.86}(\omega_{B_c})^{+0.81}_{-0.74}(f_{B_c})^{+0.35}_{-0.32}(m_c)^{+0.37}_{-0.35}(C_D)^{+0.33}_{-0.32}(f_D) \pm 0.34(\tau_{B_c})) \cdot 10^{-9}$
$Br(B_c^- \to D^- \tau^+ \tau^-)$	$(1.03^{+0.38}_{-0.27}(\omega_{B_c})^{+0.22}_{-0.20}(f_{B_c})^{+0.12}_{-0.10}(m_c)^{+0.09}_{-0.08}(C_D) \pm 0.09(f_D) \pm 0.09(\tau_{B_c})) \cdot 10^{-9}$
$Br(B_c^- \to D^- \bar{\nu}\nu)$	$(3.13^{+0.96}_{-0.71}(\omega_{B_c})^{+0.67}_{-0.61}(f_{B_c})^{+0.30}_{-0.26}(m_c)^{+0.31}_{-0.29}(C_D)^{+0.28}_{-0.26}(f_D) \pm 0.28(\tau_{B_c})) \cdot 10^{-8}$
$Br(B_c^- \to D_s^- l^+ l^-)$	$(1.56^{+0.46}_{-0.36}(\omega_{B_c})^{+0.33}_{-0.30}(f_{B_c})^{+0.17}_{-0.15}(m_c)^{+0.13}_{-0.12}(C_{D_s}) \pm 0.07(f_{D_s}) \pm 0.14(\tau_{B_c})) \cdot 10^{-7}$
$Br(B_c^- \to D_s^- \tau^+ \tau^-)$	$(0.38^{+0.13}_{-0.10}(\omega_{B_c})^{+0.08}_{-0.07}(f_{B_c})^{+0.05}_{-0.04}(m_c) \pm 0.03(C_{D_s}) \pm 0.02(f_{D_s}) \pm 0.03(\tau_{B_c})) \cdot 10^{-7}$
$Br(B_c^- \to D_s^- \bar{\nu}\nu)$	$(1.29^{+0.39}_{-0.30}(\omega_{B_c})^{+0.28}_{-0.25}(f_{B_c})^{+0.14}_{-0.12}(m_c)^{+0.11}_{-0.10}(C_{D_s}) \pm 0.06(f_{D_s}) \pm 0.11(\tau_{B_c})) \cdot 10^{-6}$
$Br(B_c^- \to D^{*-} l^+ l^-)$	$(1.21^{+0.36}_{-0.28}(\omega_{B_c})^{+0.26}_{-0.23}(f_{B_c})^{+0.14}_{-0.12}(m_c) \pm 0.11(C_{D^*})^{+0.25}_{-0.23}(f_{D^*}) \pm 0.11(\tau_{B_c})) \cdot 10^{-8}$
$Br(B_c^- \to D^{*-} \tau^+ \tau^-)$	$(0.16^{+0.05}_{-0.04}(\omega_{B_c}) \pm 0.03(f_{B_c}) \pm 0.02(m_c) \pm 0.01(C_{D^*})^{\pm} 0.03(f_{D^*}) \pm 0.01(\tau_{B_c})) \cdot 10^{-8}$
$Br(B_c^- \to D^{*-} \bar{\nu}\nu)$	$(1.10^{+0.34}_{-0.26}(\omega_{B_c})^{+0.24}_{-0.21}(f_{B_c})^{+0.13}_{-0.11}(m_c) \pm 0.10(C_{D^*})^{+0.23}_{-0.21}(f_{D^*}) \pm 0.10(\tau_{B_c})) \cdot 10^{-7}$
$Br(B_c^- \to D_s^{*-} l^+ l^-)$	$(4.40^{+1.40}_{-1.05}(\omega_{B_c})^{+0.95}_{-0.85}(f_{B_c})^{+0.72}_{-0.57}(m_c)^{+0.32}_{-0.31}(C_{D_s^*})^{+0.92}_{-0.84}(f_{D_s^*}) \pm 0.39(\tau_{B_c})) \cdot 10^{-7}$
$Br(B_c^- \to D_s^{*-} \tau^+ \tau^-)$	$(0.52^{+0.18}_{-0.13}(\omega_{B_c})^{+0.11}_{-0.10}(f_{B_c})^{+0.10}_{-0.08}(m_c) \pm 0.03(C_{D_s^*})^{+0.11}_{-0.10}(f_{D_s^*}) \pm 0.05(\tau_{B_c})) \cdot 10^{-7}$
$Br(B_c^- \to D_s^{*-} \bar{\nu}\nu)$	$(4.04^{+1.30}_{-0.97}(\omega_{B_c})^{+0.87}_{-0.78}(f_{B_c})^{+0.68}_{-0.53}(m_c)^{+0.29}_{-0.28}(C_{D_s^*})^{+0.85}_{-0.77}(f_{D_s^*}) \pm 0.36(\tau_{B_c})) \cdot 10^{-6}$

and in Table IV, we have the following points:

(i) All the form factors for the transitions $B_c \to D_s^{(*)}$ are larger than the corresponding values for the transitions $B_c \to D^{(*)}$ at $q^2 = 0$, which characterizes the SU(3) breaking effect.

(ii) $F_0(0)$ equals to $F_+(0)$ by definition for the $B_c \to D$ or $B_c \to D_s$ transition, but they have different q^2 dependence by the different parameters (a, b). $T_1(0)$ equals to $T_2(0)$ for the $B_c \to D^*$ or $B_c \to D_s^*$ transition claimed by the Eq. (17) as they are given in Table II although their expressions are different in Eqs.(25, 26).

(iii) Because of the phase space suppression, the branching ratios of the decay modes with a τ in the final product are smaller than those decay modes with electron or muon in the final product for the the charged current process. And for the flavor changing neutral current processes, with two τ's in the final product, the branching ratios are much smaller than the corresponding decays with electron or muon pairs in the final product.

(iv) The branching ratios of the decay modes with $\bar{\nu}\nu$ are almost an order magnitude larger than the corresponding decays with $l^+ l^-$ after the summation over the three neutrino generations. Because of the strong suppression of the CKM factor $|V_{td}/V_{ts}|^2 = |0.211|^2$ [24], the branching ratios for the decay modes with $b \to d$ transitions are much smaller than those decay modes with the $b \to s$ transitions.

In order to reduce the theoretical uncertainty of the form factor calculations, we define

two ratios R_D and R_{D^*} among the branching ratios for the the charged-current processes

$$R_D = \frac{Br(B_c^- \to \bar{D}^0 \tau^- \bar{\nu}_\tau)}{Br(B_c^- \to \bar{D}^0 l^- \nu_l)} = 0.69 \pm 0.01(\omega_{B_c})^{+0.01}_{-0.00}(m_c) \,, \tag{35}$$

$$R_{D^*} = \frac{Br(B_c^- \to \bar{D}^{*0} \tau^- \bar{\nu}_\tau)}{Br(B_c^- \to \bar{D}^{*0} l^- \nu_l)} = 0.59^{+0.00}_{-0.01}(\omega_{B_c})^{+0.00}_{-0.01}(m_c) \,, \tag{36}$$

with $l = (e, \mu)$. These two relations will be tested by experiments.

V. SUMMARY

In this paper we studed the $B_c \to (D_{(s)}, D^*_{(s)})$ transition form factors $F_{0,+,T}(q^2)$ and $V(q^2), A_{0,1,2}(q^2), T_{1,2,3}(q^2)$ in the pQCD factorization approach based on k_T factorization. The pQCD predictions for the values of the $B_c \to D_{(s)}$ and $B_c \to D^*_{(s)}$ transition form factors agree with those obtained using other methods. Utilizing these form factors, we calculated the branching ratios for all the semileptonic decays of $B_c^+ \to D^{(*)}_{(s)}(l^+\nu_l, l^+l^-, \nu\bar{\nu})$. Because of phase space suppression, the production ratios of the decay modes with lepton τ in the final product are smaller than the corresponding decays with electron or muon in the final product. The branching ratios of the decay modes with $\bar{\nu}\nu$ are almost an order magnitude larger than the corresponding decays with l^+l^- after the summation over the three neutrino generations. The branching ratios for the decays with the $b \to d$ transition are much smaller than those with the $b \to s$ transitions.

In order to reduce the theoretical uncertainty of the pQCD predictions, we defined two ratios R_D and R_{D^*} among the branching ratios for the the charged-current processes. The pQCD predictions are

$$R_D = \frac{Br(B_c^- \to \bar{D}^0 \tau^- \bar{\nu}_\tau)}{Br(B_c^- \to \bar{D}^0 l^- \nu_l)} \approx 0.7 \,, \tag{37}$$

$$R_{D^*} = \frac{Br(B_c^- \to \bar{D}^{*0} \tau^- \bar{\nu}_\tau)}{Br(B_c^- \to \bar{D}^{*0} l^- \nu_l)} \approx 0.6 \,, \tag{38}$$

with $l = (e, \mu)$. It would possible to test these predictions by LHCb and the forthcoming Super-B experiments.

Acknowledgments

This work is supported in part by the National Natural Science Foundation of China under Grant No. 11375208, 11228512, and 11235005.

[1] I.P. Gouz, V.V. Kiselev, A.K. Likhoded, V.I. Romanovsky, and O.P. Yushchenko, Phys. At. Nucl. **67**, 1559 (2004).

[2] N. Brambilla *et al.,* (Quarkonium Working Group), Report No. CERN-2005-005; M.P. Altarelli and F. Teubert, Int. J. Mod. Phys. A **23**, 5117 (2008).

[3] R. Dhir and R.C. Verma, Phys. Scr. **82**, 065101 (2010).
[4] A. Faessler, Th. Gutsche, M.A. Ivanov, J.G. Körner, and V.E. Lyubovitskij, Eur. Phys. J. direct C **4**, 1 (2002).
[5] C.Q. Geng, C.W. Hwang, and C.C. Liu, Phys. Rev. D **65**, 094037 (2002).
[6] D. Ebert, R.N. Faustov, and V.O. Galkin, Phys. Rev. D **82**, 034032 (2010).
[7] I. Ahmed, M. A. Paracha, M. Junaid, A. Ahmed, A. Rehman and M. J. Aslam, arXiv:1107.5694.
[8] A. Ahmed, I. Ahmed, M. A. Paracha, M. Junaid, A. Rehman and M. J. Aslam, arXiv:1108.1058.
[9] K. Azizi, F. Falahati, V. Bashiry, and S.M. Zebarjad, Phys. Rev. D **77**, 114024 (2008).
[10] K. Azizi and R. Khosravi, Phys. Rev. D **78**, 036005 (2008).
[11] T.W. Yeh and H.-n. Li, Phys. Rev. D **56**, 1615 (1997); H.-n. Li and H.L. Yu, Phys. Rev. Lett. **74**, 4388 (1995); Phys. Lett. B **353**, 301 (1995); Phys. Rev. D **53**, 2480 (1996).
[12] Y.Y. Keum, H.-n. Li, and A.I. Sanda, Phys Lett. B **504**, 6 (2001); Phys. Rev. D **63**, 054008 (2001); C. -D. Lu, K. Ukai and M. -Z. Yang, Phys. Rev. D **63**, 074009 (2001); C. -D. Lu and M. -Z. Yang, Eur. Phys. J. C **23**, 275 (2002).
[13] G. Sterman, Phys. Lett. B **179**, 281 (1986); Nucl. Phys. **B281**, 310 (1987); S. Catani and L. Trentadue, Nucl. Phys. **B327**, 323 (1989); H.-n. Li, Phys. Rev. D **66**, 094010 (2002).
[14] W.F. Wang and Z.J. Xiao, Phys. Rev. D **86**, 114025 (2012).
[15] W.F. Wang, Y.Y. Fan, M. Liu, and Z.J. Xiao Phys. Rev. D **87**, 097501 (2013).
[16] R.H. Li, C.D. Lü, and W. Wang Phys. Rev. D **79**, 034014 (2009); R.H. Li, C.D. Lü, W. Wang, and X.X. Wang, Phys. Rev. D **79**, 014013 (2009);
[17] Z.J. Xiao and X. Liu, Phys. Rev. D **84**, 074033 (2011); X. Liu, Z.J. Xiao, and C.D. Lü , Phys. Rev. D **81**, 014022 (2010); J.F. Cheng, D.S. Du, and C.D. Lü, Eur. Phys. J. C **45**,711 (2006); J.F. Sun, D.S. Du, and Y.L. Yang, Eur. Phys. J. C **60**, 107 (2009).
[18] W.F. Wang, Y.Y. Fan, and Z.J. Xiao, Chin. Phys. C **37**, 093102 (2013).
[19] T. Kurimoto, H.-n. Li, and A.I. Sanda, Phys. Rev. D **67**, 054028 (2003).
[20] J.F. Cheng, D.S. Du, and C.D. Lü, Eur. Phys. J. C **45**, 711 (2006).
[21] X. Liu, Z.J. xiao, and C.D. Lü, Phys. Rev. D **81**, 014022 (2010).
[22] Z.T. Zou, X. Yu, and C.D. Lü, Phys. Rev. D **87**, 074027 (2013).
[23] R.H. Li, C.D. Lü, and H. Zou, Phys. Rev. D **78**, 014018 (2008).
[24] J. Beringer, *et al.* (Particle Data Group), Phys. Rev. D **86**, 010001 (2012).
[25] D. Becirevic, Ph. Boucaud, J.P. Leroy, V. Lubicz, G. Martinelli, F. Mescia, and F. Rapuano, Phys. Rev. D **60**, 074501 (1999); M.A. Ivanov, J.G. Korner, P. Santorelli, Phys. Rev. D **73**, 054024 (2006); G.L. Wang, Phys. Lett. B **633**, 492 (2006); D. Becirevic, V. Lubicz, F. Sanfilippo, S. Simula, and C. Tarantino, J. High Energy Phys. 02 (2012) 042; P. Gelhausen, A. Khodjamirian, A.A. Pivovarov, and D. Rosenthal, Phys. Rev. D **88**, 014015 (2013);
[26] M. Wirbel, B. Stech, and M. Bauer, Z. Phys. C **29**, 637 (1985).
[27] M. Beneke and Th. Feldmann, Nucl. Phys. **B592**, 3 (2001).
[28] C.D. Lü and M.Z. Yang, Eur. Phys. J. C **28**, 515 (2003).
[29] T. Kurimoto, H.-n. Li, and A. I. Sanda, Phys. Rev. D **65**, 014007 (2001).
[30] P. Colangelo, F. De Fazio, P. Santorelli, and E. Scrimieri, Phys. Rev. D **53**, 3672 (1996).
[31] H.-n. Li, Y.L. Shen, and Y.M. Wang, Phys. Rev. D **85**, 074004 (2012).
[32] W. Wang, Y.L. Shen, and C.D. Lü, Phys. Rev. D **79**, 054012 (2009).
[33] A. J. Buras, M. Misiak, M. Munz, and S. Pokorski, Nucl. Phys. **B424**, 374 (1994);
G. Buchalla, A. J. Buras, and M. E. Lautenbacher, Rev. Mod. Phys. **68**, 1125 (1996).

[34] M. Beneke, Th. Feldmann, and D. Seidel, Nucl. Phys. **B612**, 25 (2001).
[35] W.J. Li, Y.B. Dai, and C.S. Huang, Eur. Phys. J. C **40**, 565 (2005).
[36] A. Ali, P. Ball, L. T. Handoko, and G. Hiller, Phys. Rev. D **61**, 074024 (2000).
[37] T. M. Aliev, A. S. Cornell, and Naveen Gaur, J. High Energy Phys. 07 (2007) 072; T. Barakat, New J. Phys. **4**, 25 (2002).
[38] P. del Amo Sanchez *et al.* (BABAR Collaboration), Phys. Rev. D **83**, 032007 (2011); Phys. Rev. D **83**, 052011 (2011).
[39] T.W. Chiu, T.H. Hsieh, C.H. Huang, and K. Ogawa (TWQCD Collaboration), Phys. Lett. B **651**, 171 (2007).
[40] D. Ebert, R. N. Faustov, and V. O. Galkin, Phys. Rev. D **68**, 094020 (2003).
[41] M. A. Ivanov, J. G. Körner, and O, N, Pakhomova, Phys. Lett. B **555**, 189 (2003).
[42] T. Huang, and F. Zuo, Eur. Phys. J. C **51**, 833 (2007).
[43] V.V. Kiselev, arXiv:hep-ph/0211021.
[44] R. Dhir and R.C. Verma, Phys. Rev. D **79**, 034004 (2009).